

NURTIVO® IRON

NURTIVO® IRON is a unique formulation of nutrient-infused amino acid iron. It has a unique package of biostimulants and organic acids for the purpose of correcting and preventing iron deficiencies and enhancing a plant's ability in the functions of chlorophyll synthesis and if left unregulated, causes a lack of chlorophyll and makes the leaves turn yellow and the growth stunt and iron chlorosis.

NURTIVO® IRON has a biostimulant package containing specific amino acids, natural plant hormones and various other organic acids and plant extracts all which aid in the plant's ability to carry out these functions to avoid iron chlorosis.

ANALYSIS 4% IRON w/v

NURTIVO® IRON is infused with multiple amino and organic acid compounds along with biostimulants to amplify the nutrient uptake and metabolic activity.

APPLICATION RATES

GENERAL RATES INFORMATION: This is a foliar-applied nutritional supplement used in agricultural applications that must be used diluted in water. When used as directed it is nontoxic to plant species specified.

Application rates should always be declared after having an agronomist or agricultural consultant take into consideration the entire fertiliser program and other nutrients being applied at the same time as this product. As some fungicides containing copper and sulphur can cause phytotoxicity issues when combined with or used near the foliar fertiliser application all care should be taken, and if combined with other chemistry, seek professional advice from your NURTIVO® representative.

Rates have been determined based on full-sized plants. If foliar applications are to be made on new plantations or immature plants then an agronomist or agricultural consultant should advise a lower usage rate based on the situation.

TREE CROPS: Apply at rates of 2.0–5.0L per hectare once tree growth has commenced. Multiple applications can be made throughout the season when the requirement for iron is required. When using to correct iron deficiencies or disorders use the higher rate once or twice in succession 5–10 days apart or until the deficiency has been corrected.

NURTIVO® IRON APPLICATIONS:



GRAPES



ALMONDS



CITRUS



VEGETABLES



POTATOES



ONIONS

GRAPEVINES: Apply at rates of 2.5–4.0L per hectare once vine growth has commenced. Multiple applications can be made throughout the season when the requirement for iron is required. When using to correct iron deficiencies or disorders use the higher rate once or twice in succession 5–10 days apart or until the deficiency has been corrected.

BERRIES: Apply at rates of 1.0–3.0L per hectare once growth has commenced. Multiple applications can be made throughout the season when the requirement for iron is required. When using to correct iron deficiencies or disorders use the higher rate once or twice in succession 5–10 days apart or until the deficiency has been corrected.

FIELD CROPS & VEGETABLES: Apply at 1.0–2.0L per hectare once growth has commenced. Multiple applications can be made throughout the season when the requirement for iron is required. When using to correct iron deficiencies or disorders use the higher rate once or twice in succession 5–10 days apart or until the deficiency has been corrected.

BENEFITS OF NURTIVO® IRON:



Leaf Health



Controlled
Vegetative Growth



Stem Health



Disease Resistance