

# NURTIVO<sup>®</sup>

## ZINC

**NURTIVO<sup>®</sup> ZINC** is a unique formulation of nutrient-infused amino acid zinc.

It has a unique package of biostimulants and organic acids for the purpose of correcting and preventing zinc deficiencies and enhancing a plant's ability in the functions of chlorophyll formation, certain carbohydrates formation, and leaf health.

Zinc is essential in the formation of auxins which help with growth regulation and stem elongation. It has a biostimulant package containing specific amino acids, natural plant hormones and various other organic acids and plant extracts all of which aid in the plant's ability to carry out these functions.

### NURTIVO<sup>®</sup> ZINC APPLICATIONS:



GRAPES



ALMONDS



CITRUS



VEGETABLES



POTATOES



ONIONS

### ANALYSIS

**5% ZINC w/v**

**NURTIVO<sup>®</sup> ZINC** is infused with multiple amino and organic acid compounds along with biostimulants to amplify the nutrient uptake and metabolic activity.

### APPLICATION RATES

**GENERAL RATES INFORMATION:** This is a foliar-applied nutritional supplement used in agricultural applications that must be used diluted in water. When used as directed it is nontoxic to plant species specified.

Application rates should always be declared after having an agronomist or agricultural consultant take into consideration the entire fertiliser program and other nutrients being applied at the same time as this product. As some fungicides containing copper and sulphur can cause phytotoxicity issues when combined with or used near the foliar fertiliser application all care should be taken, and if combined with other chemistry, seek professional advice from your NURTIVO<sup>®</sup> representative.

Rates have been determined based on full-sized plants. If foliar applications are to be made on new plantations or immature plants then an agronomist or agricultural consultant should advise a lower usage rate based on the situation.

**TREE CROPS:** Apply at rates of 2.0–4.0L per hectare once tree growth has commenced. Multiple applications can be made throughout the season when the requirement for zinc is required. When using to correct zinc deficiencies or disorders use the higher rate once or twice in succession 5–10 days apart or until the deficiency has been corrected.

**GRAPEVINES:** Apply at rates of 2.5–4.0L per hectare once vine growth has commenced. Multiple applications can be made throughout the season when the requirement for zinc is required. When using to correct zinc deficiencies or disorders use the higher rate once or twice in succession 5–10 days apart or until the deficiency has been corrected.

**BERRIES:** Apply at rates of 1.0–3.0L per hectare once growth has commenced. Multiple applications can be made throughout the season when the requirement for zinc is required. When using to correct zinc deficiencies or disorders use the higher rate once or twice in succession 5–10 days apart or until the deficiency has been corrected.

**FIELD CROPS & VEGETABLES:** Apply at 1.0–2.0L per hectare once growth has commenced. Multiple applications can be made throughout the season when the requirement for zinc is required. When using to correct zinc deficiencies or disorders use the higher rate once or twice in succession 5–10 days apart or until the deficiency has been corrected.

### BENEFITS OF NURTIVO<sup>®</sup> ZINC:



Stress Relief



Cell Division



Pollen Germination



Flowering



Fruit Setting



Skin Walls